

First record of Norwegian killer whales attacking and feeding on a harbour porpoise

A. MEL COSENTINO^{1,2}

¹Institute of Biological and Environmental Sciences, University of Aberdeen, School of Biological Sciences, Tillydrone Avenue, Aberdeen, AB24 2TZ, UK, ²Hvalsafari A.S., Hamnegata 1B, 8480 Andenes, Norway

Orcinus orca is a cosmopolitan species and the most widely distributed marine mammal. Its diet includes over 140 species of fish, cephalopods, sea birds and marine mammals. However, many populations are specialised on certain specific prey items. Three genetically distinct populations have been described in the North Atlantic. Population A (that includes the Icelandic and Norwegian sub-populations) is believed to be piscivorous, as is population C, which includes fish-eating killer whales from the Strait of Gibraltar. In contrast, population B feeds on both fish and marine mammals. Norwegian killer whales follow the Norwegian spring spawning herring stock. The only description in the literature of Norwegian killer whales feeding on another cetacean species is a predation event on northern bottlenose whales in 1968. Daily land-based surveys targeting sperm whales were conducted from the Andenes lighthouse using BigEyes[®] binoculars (25×, 80 mm). The location of animals at sea was approximated through the use of an internal reticule system and a graduated wheel. On 24 June 2012 at 3:12 am, an opportunistic sighting of 11 killer whales was made off Andenes harbour. The whales hunted and fed on a harbour porpoise. Despite these species having overlapping distributions in Norwegian waters, this is the first predatory event reported in the literature.

Keywords: killer whale, diet, behaviour, harbour porpoise

Submitted 8 April 2015; accepted 18 June 2015

INTRODUCTION

The killer whale (*Orcinus orca*, Linnaeus 1758) is a cosmopolitan species and the most widely distributed marine mammal. It can be found in all of the world's oceans, from the equator to polar waters (Forney & Wade, 2007), and shows seasonal movement patterns that are usually associated with increased prey abundance (López & López, 1985; Similä *et al.*, 1996; Ford *et al.*, 1998; Iñiguez, 2001; Esteban *et al.*, 2013). Worldwide, its diet includes over 140 species of fish, cephalopods, sea birds, turtles and marine mammals. However, most studied killer whale populations are specialized to feed on certain specific prey items (Baird, 1994; Ford *et al.*, 1998; Pitman & Ensor, 2003; Herman *et al.*, 2005; Pitman *et al.*, 2007; Esteban, 2008).

The best studied killer whale populations in the world are found off the west coast of the US and Canada, where three different ecotypes have been described: a 'transient' type that specializes in feeding on marine mammals and sea birds; a 'resident' type that feeds on fish in nearshore waters and a piscivorous 'offshore' type (Bigg *et al.*, 1990; Ford *et al.*, 1998; Herman *et al.*, 2005). These ecotypes not only differ in their diet, but also in their morphological traits, acoustic behaviour and social structure. Transient and resident ecotypes killer whales have also been documented off the Kamchatka peninsula and in the Sea of Okhotsk, in Russian waters (Burdin *et al.*, 2004).

Using microsatellites and mtDNA, three genetically distinct populations have been identified in the North Atlantic (Foote *et al.*, 2011). A piscivorous population A specialized in feeding on herring (*Clupea harengus*), the Norwegian, the Icelandic or the North Sea herring stocks. In turn, this population is divided into two sub-populations: the Icelandic and the Norwegian (Foote *et al.*, 2012). Individuals from population A were all sampled at latitudes above 60°N. Killer whales from population B feed on both fish and marine mammals, and live in sympatry with population A in part of their range, as individuals were sampled between latitudes 51°N and 66°N, and from the North Sea to the West coast of Iceland (Foote *et al.*, 2011). Population C includes all individuals sampled in the Strait of Gibraltar, which are specialized in feeding on bluefin tuna (*Thunnus thynnus*) (Esteban, 2008), and individuals sampled in the Canary Islands (Foote *et al.*, 2011), whose feeding habits are unknown.

Although killer whales have been recorded along the entire coast of Norway, as well as in offshore waters (Foote *et al.*, 2007), most of what is known about Norwegian killer whales comes from research studies carried out in northern Norway, especially during the winter time. These whales have been intensively studied since 1983 in the Vestfjord–Tysfjord–Ofotfjord area (Lofoten Archipelago), where they were observed to return every year to feed on the over-wintering Norwegian spring-spawning herring (NSSH) (Similä *et al.*, 1996; Kuningas *et al.*, 2007). Few killer whale observations are made during the summer months, but they have been recorded both off Andenes (Vesterålen Archipelago) and the Lofoten Islands (Similä *et al.*, 1996; Similä, 1997; Ugarte, 2001; Stenersen & Similä, 2004). These whales show a

Corresponding author:

A. Mel Cosentino

Email: orcinus.orca.1758@gmail.com

preference for herring, and photo-identification and satellite tags suggest that at least some groups follow the NSSH stock, year round (Similä *et al.*, 1996; Stenersen & Similä, 2004; Foote *et al.*, 2011). In addition, they have also been observed feeding on mackerel (*Scomber scombrus*), cod (*Gadus morhua*, Linnaeus 1758), Atlantic salmon (*Salmo salar*, Linnaeus 1758) and saithe (*Pollachius virens*, Linnaeus 1758), as well as harbour seals (*Phoca vitulina*, Linnaeus 1758) and, on rare occasions, sea birds (Similä *et al.*, 1996; Stenersen & Similä, 2004; Vester & Hammerschmidt, 2013).

Worldwide, there have been reports of killer whale predation on more than 20 other cetacean species, ranging from small porpoises to great whales (e.g. Hoyt, 1990; Jefferson *et al.*, 1991). However, for the Norwegian population, there is only a report on a single predation event on northern bottlenose whales (*Hyperoodon ampullatus*, Forster 1770) (Jonsgård, 1968 in Similä *et al.*, 1996). Here I report and describe the first case in which Norwegian killer whales have been observed hunting and feeding on a harbour porpoise (*Phocoena phocoena*, Linnaeus 1758).

MATERIALS AND METHODS

The event reported here occurred off Andenes harbour in northern Norway, at approximately 69.334°N 16.161°E. Since 2011, several land-based surveys were carried out in the area on a daily basis (depending on weather conditions), as part of a study focusing on sperm whales (*Physeter macrocephalus*, Linnaeus 1758). Such surveys were conducted from the Andenes lighthouse using BigEyes® binoculars (25× magnifications, 80 mm). The lighthouse is located at the northernmost point of Andøy Island (69.3240°N 16.1159°E) and the binoculars are assembled on the gallery deck at approximately 40 meters above sea level. For the above-mentioned sperm whale surveys, the study area is divided into two contiguous areas (each 120° wide): Bleik Canyon on the west side and Andfjord on the east side of the island (Figure 1). Surveys involve scanning one of the two areas for 2 out of every 5 min, for a period of 1 h (i.e. 12 scans per hour). Data are

collected using an Olympus Recorder WS-750M. Location of animals at sea is approximated through the use of an internal reticule system and a graduated wheel.

The 24 h of daylight during the summer months in the study area makes it possible to conduct land-based surveys at any time of day.

RESULTS

On 24 June 2012 at 03:10, an opportunistic sighting of a group of 11 killer whales was made off Andenes in Andfjord waters (69.28486°N 16.18732°E) (Figure 2). Based on body size, as well as size and shape of the dorsal fin, it was determined that the group was composed of four adult males, one calf, one juvenile and five subadult individuals/adult females. The whales were slowly travelling northwards in a loose formation, less than two body lengths from each other (*sensu* Barrett-Lennard *et al.*, 1996). After 36 min, during a new scan, the group was spotted again, still travelling northwards and in a loose formation. At 03:55 they were seen less than 2000 m north-east off Andenes harbour, in a 20 m deep reef area (69.32942°N 16.17670°E). The group was heading north-east when, suddenly, the entire group turned southwards and started porpoising at high speed towards the shore. It was then decided to discontinue the sperm whale survey and perform a focal follow on the killer whales instead. Within one minute, a killer whale (a subadult or an adult female) rammed a harbour porpoise from below forcing it out of the water and exposing half of its own body at an approximately 75° angle. The porpoise was lifted into the air about 5 m above the killer whale. High-speed chasing above the water is commonly observed during killer whale predation on small- and medium-sized cetaceans in other parts of the world (e.g. Baird, 1994; Constantine *et al.*, 1998; Visser *et al.*, 2010; Coscarella *et al.*, 2015); however, this was not the case on this occasion, given that the porpoise was first observed when rammed in the air. The rest of the group continued porpoising for a few seconds, gathering around the spot where both the killer whale and the porpoise were last seen. After that, for




Fig. 1. Map of the study area in northern Norway.




Fig. 2. Map showing the locations where the killer whales were observed.

less than 1 min, the entire group remained underwater. Suddenly a whale (again, a subadult or an adult female) rose out the water as previously, but this time carrying the porpoise in its mouth. Once more, the group gathered around the attacking individual and started swimming erratically in the area. Within the next few minutes, the porpoise was rammed in the air once more, though not as high as the first time. The porpoise was not seen again and the killer whales came closer to each other, moving slowly, gathering around a small area and facing towards where the porpoise was last observed, suggesting prey consumption. Reuniting around the prey during handling and after the kill suggests prey sharing (e.g. Hoelzel, 1991; Baird, 1994). The event took place in just over 6 min. The group remained in the area and sea birds slowly gathered around, flying in circles above the killer whales, occasionally plunging around the animals.

Whilst the adult males were not actively involved and stayed to the side most of the time, the calf remained close to the group throughout the whole predation event. This has also been observed in North Pacific transients (Baird, 1994). After the kill, aerial behaviour was observed, including several breaches and tail slapping, as well as other displays such as rolling on their bodies and spyhopping, which are indicative of socialization (Barrett-Lennard *et al.*, 1996).

The whales started slowly moving to the north-east 10 min after the kill, and divided into two smaller groups after another 10 min: one male, the calf and three other individuals remained together in that spot and the rest dispersed and moved further northwards. Both groups were accompanied by birds occasionally diving, suggesting the prey was shared by both subgroups. At 04:45 all five remaining killer whales started to disperse and also headed northwards. Observation was then finished due to intense glare.

Harbour porpoises are known to be part of the diet of transient killer whales' diets off the west coast of British Columbia, Canada (Jefferson *et al.*, 1991; Baird & Guenther, 1995). Despite the fact that the species have overlapping distributions in Norwegian waters (Bjørge & Øien, 1995; Foote *et al.*, 2007), no predatory interaction has been described previously in the scientific literature. In April 2012 a photograph taken by a tourist in Eikelandsfjorden in Hardanger, Southern Norway, over 1500 km from our study area, was published in an online local newspaper,¹ showing a killer whale (a subadult

or an adult female) ramming a harbour porpoise. However, it is not known whether the porpoise was killed and consumed by the whales.

It was, unfortunately, impossible to take photographs using the binoculars from the land station for photo-identification. However, the short distance to the event made it possible to make reliable sketches of the dorsal fins of three individuals that had distinctive nicks: an adult male, a juvenile and a third individual, most likely to be an adult female. The sketches and group composition strongly suggest that the group was the same as one seen near Stø on 21 June 2012 (three days before), less than 30 km south-west of Andenes (69.17499°N 15.39300°E). Photographs taken on that occasion identified a total of 11 individuals, including one calf, one juvenile, and four adult males. It was impossible to attempt to photographically identify the whale in the photograph published in the online newspaper due to the low quality of the available image.

DISCUSSION

The above observations can be explained in one of three ways. First, this may indicate an expansion in the range of mammal-eating population B. Our current knowledge on their distribution range is limited and while a range increase in association with climate change (e.g. MacLeod, 2009) or other factors should not be disregarded, this situation is probably unlikely given the distance between the study area and the known range of population B. The other two possible explanations are that members of population A may have learned to take new prey animals only recently due to a changing environment, or that this behaviour has simply gone unnoticed to date. Both are likely situations given the already reported opportunistic foraging behaviour of this population, and that most of what is known about the diet and feeding behaviour of Norwegian killer comes from studies carried out in the wintering grounds of the NSSH stock (Similä *et al.*, 1996; Similä, 1997; Ugarte, 2001; Stenersen & Similä, 2004; Kuningas *et al.*, 2007), despite the fact that several groups have been observed in different seasons and study areas.

Recent studies suggest that the behavioural ecology of killer whales off Scotland and Iceland is more consistent with specialization in foraging behaviour at the individual or group level rather than population level (Beck *et al.*, 2011). The possibility that the Norwegian whales also show group or individual specialization in foraging behaviour cannot be disregarded. In fact, Similä *et al.* (1996) already noticed that at least some groups had different prey preferences (e.g. young herring instead of adults) associated with different seasonal occurrences. More recently, killer whales have been regularly observed feeding on harbour seals off Stø (northern tip of the island of Langøya) during the summer months (C. Ilmoni, personal communication) and on salmon in the Lofoten archipelago (Vester & Hammerschmidt, 2013), although it is not yet known if different groups are engaged in these activities.

In addition to the porpoise event, I recorded two observations of killer whales in what appeared to be foraging behaviour in search of seals on rocky islands in nearshore waters (as described in Beck *et al.*, 2011). The first sighting occurred on 2 August 2012 off Stø, when a group of five killer whales, initially observed feeding on fish (possibly mackerel), switched to actively searching for seals around the rocky islands in the

¹<http://www.nrk.no/hordaland/fekk-blinkskot-av-spekkhoggar-1.8065404>

area (Beck *et al.*, 2011). On this occasion, at least one unsuccessful attempt to catch a seal was made. Similar behaviour was observed by a group of 20 to 25 whales on the 28 November 2012, off Andenes (on the west side of Andøy Island), although no attempt to catch a seal was observed. Interestingly, one subgroup in this second sighting caught (and presumably ate) several sea birds (possibly King Eider – *Somateria spectabilis* Linnaeus 1758). There are also reports of killer whales harassing and probably feeding on puffins (*Fratercula arctica* Linnaeus 1758) in the area (G. Mann, personal communication). Together, these observations may indicate that the killer whales in northern Norway prey upon an array of fish, sea bird and marine mammal species, even though its presence in the area is highly linked to the occurrence of the NSSH (Similä *et al.*, 1996; Hvalsafari unpublished data). Furthermore, observations of killer whales harassing other cetacean species (e.g. minke and sperm whales) off Andenes have been made since 2011 (Hvalsafari, unpublished data). However, it is not clear whether or not they were predatory interactions.

The ecological separation of the North Atlantic killer whales is not as clearly defined as it is in the Eastern North Pacific (de Bruyn *et al.*, 2013). The NSSH has a complex life cycle that requires suitable areas for each stage (i.e. spawning, feeding, nursing and wintering). These areas and their migrations routes vary somewhat unpredictably (Dickson & Østerhus, 2007), which could explain the more generalist behaviour observed for this killer whale population.

Baird & Dill (1996) estimated that the optimal group size that maximizes energy intake for the North-east Pacific transients is three individuals, which coincides with the typical group size (Baird, 1994). Other marine-mammal-eating killer whales also have small group sizes: three individuals for the Punta Norte (Patagonia) population (Hoelzel, 1991), and five for killer whales in Scottish waters (Beck *et al.*, 2011). Group size for Norwegian killer whales, however, ranges from 6 to 30 animals, with a median of 15 (Similä, 1997). Foraging techniques are transferred from one generation to the other through social learning, as observed in Patagonia where killer whales use an intentional stranding technique to catch seal pups on the beach (López & López, 1985; Hoelzel, 1991).

As a population, Norwegian killer whales feed on various prey species, ranging from fish, to birds to other marine mammals, many requiring different complex hunting techniques that require a high level of coordination for a successful attack. The hunting behaviour described here greatly differs from the carousel method used when feeding on herring (Similä & Ugarte, 1993) or that used for seal hunting (Beck *et al.*, 2011). Thus, this report increases our knowledge of killer whale diet and foraging behaviour in the area and may suggest a wider separation from the Icelandic sub-population than previously thought. Progress in the knowledge of this killer whale population is slow due to the paucity of published reports. Increased collaboration between research groups in the area will greatly benefit the understanding of their feeding ecology at an individual and group level, as well as the ecology and social structure.

ACKNOWLEDGEMENTS

I would like to thank Arctic Whale Tours for providing the killer whale photographs, and two anonymous reviewers for their comments on early versions of this manuscript.

REFERENCES

- Baird R.W. (1994) *Foraging behaviour and ecology of transient killer whales* (*Orcinus orca*). PhD thesis, Simon Fraser University, Vancouver, Canada.
- Baird R.W. and Dill L.M. (1996) Ecological and social determinants of group size in transient killer whales. *Behavioral Ecology* 7, 408–416.
- Baird R.W. and Guenther T.J. (1995) Account of harbour porpoise (*Phocoena phocoena*) strandings and bycatches along the coast of British Columbia. *Reports of the International Whaling Commission, Special Issue* 16, 159–168.
- Barrett-Lennard L.G., Ford J.K.B. and Heise K.A. (1996) The mixed blessing of echolocation: differences in sonar use by fish-eating and mammal-eating killer whales. *Animal Behaviour* 51, 553–565.
- Beck S., Kuningas S., Esteban R. and Foote A.D. (2011) The influence of ecology on sociality in the killer whale (*Orcinus orca*). *Behavioral Ecology* 23, 246–253.
- Bigg M.A., Olesiuk P.F., Ellis G.M., Ford J.K.B. and Balcomb K.C. (1990) Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. *Reports of the International Whaling Commission, Special Issue* 12, 383–405.
- Bjørge A. and Øien N. (1995) Distribution and abundance of harbour porpoise, *Phocoena phocoena*, in Norwegian waters. *Reports of the International Whaling Commission, Special Issue* 16, 89–98.
- Burdin A.M., Hoyt E., Sato H., Tarasyan K.K. and Filatova O.A. (2004) Resident and transient-type killer whales, *Orcinus orca*, in Southeast Kamchatka, Russia. *International Whaling Commission Report*, SC/56/SM15, 3 pp.
- Constantine R., Visser I., Buurman D., Buurman R. and McFadden B. (1998) Killer whale (*Orcinus orca*) predation on dusky dolphins (*Lagenorhynchus obscurus*) in Kaikoura, New Zealand. *Marine Mammal Science* 14, 324–330.
- Coscarella M., Bellazzi G., Gaffet M.L., Berzano M. and Degradi M. (2015) Technique used by Killer Whales (*Orcinus orca*) when hunting for Dolphins in Patagonia, Argentina. *Aquatic Mammals* 41, 192–197.
- de Bruyn P.J.N., Tosh C.A. and Terauds A. (2013) Killer whale ecotypes: is there a global model? *Biological Reviews of the Cambridge Philosophical Society* 88, 62–80.
- Dickson B. and Østerhus S. (2007) One hundred years in the Norwegian Sea, Norsk Geografisk Tidsskrift. *Norwegian Journal of Geography* 61, 56–75.
- Esteban R. (2008) *Abundancia, estructura social y parámetros de historia natural de la orca* (*Orcinus orca*) en el Estrecho de Gibraltar. MSc thesis, Universidad de Cádiz, Cádiz, Spain. Available at: <http://www.circe.biz>
- Esteban R., Verborgh P., Gauffier P., Esteban R., Verborgh P., Gauffier P., Giménez J., Afán I., Cañadas A., García P., Murcia J.L., Magalhães S., Andreu E. and de Stephanis R. (2013) Identifying key habitat and seasonal patterns of a critically endangered population of killer whales. *Journal of the Marine Biological Association of the United Kingdom*, 1–9, doi: 10.1017/S002531541300091X.
- Foote A.D., Vester H., Víkingsson G.A. and Newton J. (2012) Dietary variation within and between populations of northeast Atlantic killer whales, *Orcinus orca*, inferred from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses. *Marine Mammal Science* 28, e472–e485, doi: 10.1111/j.1748-7692.2012.00563.x.
- Foote A.D., Víkingsson G., Øien N., Bloch D., Davis C.G., Dunn T.E., Harvey P., Mandleberg L., Whooley P. and Thompson P.M. (2007) Distribution and abundance of killer whales in the North East Atlantic.

SC/59/SM5 Report to the Scientific Committee of the International Whaling Commission, Anchorage, Alaska.

- Foote A.D., Vilstrup J.T., de Stephanis R., Verborgh P., Abel Nielsen S.C., Deaville R., Kleivane L., Vidal M., Miller P.J.O., Øien N., Pérez-Gil M., Rasmussen M., Reid R.J., Robertson K.M., Rogan E., Similä T., Tejedor M.L., Vester H., Vikingsson G.A., Willerslev E., Gilbert M.T.P. and Piertney S.B.** (2011) Genetic differentiation among North Atlantic killer whale populations. *Molecular Ecology* 20, 629–641.
- Ford J.K.B., Ellis G.M., Barrett-Lennard L.G., Morton A.B., Palm R.S. and Balcomb K.C. III** (1998) Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Canadian Journal of Zoology* 76, 1456–1471.
- Forney K.A. and Wade P.** (2007) Worldwide distribution and abundance of killer whales. In Estes J.A., DeMaster D.P., Doak D.F., Williams T.M. and Brownell R.L. Jr (eds) *Whales, whaling, and ocean ecosystems*. Berkeley: University of California Press, pp. 145–162.
- Herman D., Burrows D., Wade P., Durban J.W., Matkin C.O., LeDuc R.G., Barrett-Lennard L.G. and Krahn M.M.** (2005) Feeding ecology of eastern North Pacific killer whales *Orcinus orca* from fatty acid, stable isotope, and organochlorine analyses of blubber biopsies. *Marine Ecology Progress Series* 302, 275–291.
- Hoelzel A.R.** (1991) Killer whale predation on marine mammals at Punta Norte, Argentina: food sharing, provisioning and foraging strategy. *Behavioral Ecology and Sociobiology* 29, 197–204.
- Hoyt E.** (1990) *Orca: the whale called killer*, 3rd edn. London: Robert Hale Limited.
- Iñiguez M.A.** (2001) Seasonal distribution of killer whales (*Orcinus orca*) in northern Patagonia, Argentina. *Aquatic Mammals* 27, 154–161.
- Jefferson T.A., Stacey P.J. and Baird R.W.** (1991) A review of killer whale interactions with other marine mammals: predation to coexistence. *Mammal Review* 21, 151–180.
- Jonggård Å.** (1968) A note on the attacking behavior of the killer whale (*Orcinus orca*). *Norsk Hvalfangst-Tid* 57, 84–85.
- Kuningas S., Similä T. and Hammond P.S.** (2007) Population dynamics of killer whales (*Orcinus orca*) off northern Norway. SC/59/SM13 Report to the Scientific Committee of the International Whaling Commission, Anchorage, Alaska.
- López J.C. and López D.** (1985) Killer whales of Patagonia and their behavior of intentional stranding while hunting nearshore. *Journal of Mammalogy* 66, 181–183.
- MacLeod C.D.** (2009) Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endangered Species Research* 7, 125–136.
- Pitman R.L. and Ensor P.** (2003) Three different forms of killer whales in Antarctic waters. *Journal of Cetacean Research and Management* 5, 131–139.
- Pitman R.L., Perryman W.L., LeRol D. and Eilers E.** (2007) A dwarf form of killer whale in Antarctica. *Journal of Mammalogy* 88, 43–48.
- Similä T.** (1997) Sonar observations of killer whales (*Orcinus orca*) feeding on herring schools. *Aquatic Mammals* 23, 119–126.
- Similä T., Holst J.C. and Christensen I.** (1996) Occurrence and diet of killer whales in northern Norway: seasonal patterns relative to the distribution and abundance of Norwegian spring-spawning herring. *Canadian Journal of Fisheries and Aquatic Science* 53, 769–779.
- Similä T. and Ugarte F.** (1993) Surface and underwater observations of cooperatively feeding killer whales in northern Norway. *Canadian Journal of Zoology* 71, 1494–1499.
- Stenersen J. and Similä T.** (2004) *Norwegian killer whales*. Lofoten: Tringa Forlag, 92 pp.
- Ugarte F.** (2001) *Behaviour and social organisation of killer whales in northern Norway*. MSc thesis, University of Tromsø, Tromsø, Norway.
- Vester H. and Hammerschmidt K.** (2013) First record of killer whales (*Orcinus orca*) feeding on Atlantic salmon (*Salmo salar*) in northern Norway suggest a multi-prey feeding type. *Marine Biodiversity Records* 6, e9, doi: 10.1017/S1755267212001030.

and

Visser I.N., Zaeschmar J., Halliday J., Abraham A., Ball P., Bradley R., Daly S., Hatwell T., Johnson T., Johnson W., Kay L., Maessen T., McKay V., Peters T., Turner N., Umuroa B. and Pace D.S. (2010) First record of predation on false killer whales (*Pseudorca crassidens*) by killer whales (*Orcinus orca*). *Aquatic Mammals* 36, 195–204.

Correspondence should be addressed to:

A. Mel Cosentino
Wild Earth Foundation
Av de las Ballenas 9500
Puerto Pirámides, Península Valdes
Chubut, Argentina
email: orcinus.orca.1758@gmail.com